

## Injections, Blood Draws, and IV's



Dab don't scrub  
with alcohol wipes.



Place tourniquet over  
clothing or protective  
bandages. OR use  
manual (hands)  
tourniquet.



Do NOT use Coban.



Protect the skin with a silicone  
adhesive base layer.



Do NOT place adhesive  
dressing directly on skin.

1

## Monitors and Blood Pressure Cuff



Dab cotton ball on  
adhesive backing to  
minimize adhesive.



Hold in place  
with gauze



Do NOT put  
adhesive  
directly on  
skin




Put blood pressure  
cuff over clothing or  
gauze.

2

## Exams, Therapeutic Holds, Oral Medications



Do NOT pull on ears



Do NOT put pressure or friction on skin, including lips and internal mucosa



**Leads:** Remove backing or use cotton. Secure with mepitac or gauze.



3

## Protect the Skin with a Base Layer

**Mepiform**

Mepiform®




**Mepitac**



**Mepilex Lite**



**Mepitel/Mepitel One**



Remove BOTH sides of plastic backing if Mepitel.

**Mepilex**

Mepilex®



**Mepilex Border**



4

Protect the Skin with a Base Layer. Secure device to base layer. Use gauze for extra stability.



5

Remove all adhesives from skin slowly AND with adhesive remover



Do not pull **up** on adhesives. Fold it **back on top of itself** AND use an adhesive remover.

6

## Hospital Visits

- IV
- PICC
- Ports
- Feeding Tubes
- Leads
- Foleys
- Blood draws
- Masks
- Nasal Cannula
- BP monitors
- Transfers
- Boosting
- Hygiene
- Toileting
- Dressing Changes
- Pulse Oximeters
- Pediatric Medication Administration